

Fall 9-8-2013

Sensing of polyethylene glycols using a channel of the connector from the Phi29 bacteriophage

Peng Jing

Department of Chemistry, Indiana -Purdue University, Fort Wayne, Indiana 46805,, jingp@ipfw.edu

Rong Zhang

*Division of Endocrinology, Harvard Medical School, Children's Hospital Boston, Boston, Massachusetts, 02115,
Rong.Zhang@childrens.harvard.edu*

Follow this and additional works at: http://opus.ipfw.edu/chemistry_facpres



Part of the [Chemistry Commons](#)

Opus Citation

Peng Jing and Rong Zhang (2013). *Sensing of polyethylene glycols using a channel of the connector from the Phi29 bacteriophage*. Anyl. The American Chemical Society. Presented at 246th ACS National Meeting & Exposition, Indianapolis, IN.
http://opus.ipfw.edu/chemistry_facpres/41

This Poster Session is brought to you for free and open access by the Department of Chemistry at Opus: Research & Creativity at IPFW. It has been accepted for inclusion in Chemistry Faculty Presentations by an authorized administrator of Opus: Research & Creativity at IPFW. For more information, please contact admin@lib.ipfw.edu.

ANYL

Thomas Rossi

Sunday, September 08, 2013

69 - Sensing of polyethylene glycols using a channel of the connector from the Phi29 bacteriophage

Peng Jing¹, jingp@ipfw.edu, Rong Zhang². (1) Department of Chemistry, Indiana -Purdue University, Fort Wayne, Indiana 46805, United States, (2) Division of Endocrinology, Harvard Medical School, Children's Hospital Boston, Boston, Massachusetts 02115, United States

Nanopore sensor is a novel detection technology that uses a nano-sized pore formed either by channel proteins or by synthetic materials for detecting characteristic currents when analytes pass through it. The phi29 connector is a channel protein that *in-vivo* serves as a path for the genome DNA to enter the capsid of the bacteriophage phi29. Recently our work have shown the protein channel can be embedded into a bilayer membrane and, under an electric field, ionic current induced by ds-DNA translocation through the channel can be detected. The results suggest the channel protein would be possibly used as a central motif in the nanopore sensor.

In this study, a phi29 mutant connector containing a six histidine-tag was prepared and inserted into a bilayer membrane for nanopore sensing. A series of hydrophilic polymers, PEGs, with molecular weights ranging from 3000 Da-200000 Da are firstly used as analytes. In an identical electrolyte solution containing PEG with different molecular weights, the ionic currents induced by a single channel are recorded and compared. It was found that the single channel conductance is strongly influenced by the molecular weights of the used PEG. Since the size of a PEG molecule is determined by its molecular weight, a relationship between the resultant conductance and the size of the used PEGs has been established. The results also indicate a possibility of using the PEG to probe the internal geometry of the C-His connector channel.

Sunday, September 8, 2013 03:00 PM[General Posters \(03:00 PM - 05:00 PM\)](#)**Location: Indiana Convention Center****Room: Halls F&G**

*ACS does not own copyrights to the individual abstracts. For permission, please contact the author(s) of the abstract.

[Close Window](#)